



Poitiers
France

18.20
June
2019

© Jean Weber – Inra

International conference

Food security and climate change: 4 per mille
initiative new tangible global challenges for the soil.



BOOK OF ABSTRACTS

International conference food security climate change: 4 per mille initiative new tangible global challenges for the soil

18-20 june 2019, Poitiers - France

symposium.inra.fr/4p1000



A MODEL-BASED ASSESSMENT OF THE SOIL C STORAGE POTENTIAL AT THE NATIONAL SCALE: A CASE STUDY FROM FRANCE

Pellerin S.¹, Bamiere L.¹, Constantin J.¹, Launay C.¹, Martin R.¹, Schiavo M.¹, Angers D.², Augusto L.¹, Balesdent J.¹, Basile-Doelsch I.¹, Bellassen V.¹, Cardinael R.³, Cecillon L.⁴, Ceschia E.¹, Chenu C.⁵, Daroussin J.¹, Delacote P.¹, Delame N.¹, Gastal F.¹, Graux A.-I.¹, Guenet B.⁶, Houot S.¹, Klumpp K.¹, Letort E.¹, Martin M.¹, Mary B.¹, Menasseri S.⁷, Meziere D.¹, Mosnier C.¹, Morvan T.¹, Roger-Estrade J.⁵, Saint-Andre L.¹, Therond O.¹, Viaud V.¹, Rechauchere O.¹, Richard G.¹

¹INRA, France, ²AAFC/AAC, Canada, ³CIRAD, France, ⁴IRSTEA, France, ⁵AgroParisTech, France, ⁶CNRS, France, ⁷Agrocampus Ouest, France

The recent controversy about the 4 per 1000 initiative has emphasized the need for a quantitative assessment of the C storage potential of agricultural soils. Moreover a clear distinction is required between the biophysically and the economically achievable potentials. Here we used a modelling approach at a fine spatial-scale resolution (< 8 km²) to quantify the additional C storage in agricultural soils of mainland France following the implementation, when feasible, of a range of soil C storing practices (i.e. cover crops, reduced tillage, new C inputs, grazing instead of mowing,...). The additional cost for farmers was also calculated, thus yielding the cost per Mg of additional C stored in soils. Results showed that the average additional C storage calculated over 30 years ranged between +0.028 and +0.466 Mg C ha⁻¹ yr⁻¹ (i.e. between +0.5 and +7.2‰) for the different tested practices, with a very high spatial variability over France for each practice related to initial C stocks and pedo-climatic conditions. The storing practices were then ranked according to the cost of the additional C stored in soils (expressed in euro per Mg of C) and an optimal cost-efficient strategy was proposed at the national level.